



C-TAK1 Positive Control

Product Data Sheet

For Research Use Only, Not for use in diagnostic procedures

C-TAK1 Positive Control

(Human, full length, recombinant enzyme expressed in *E. coli*.)

Cat# CY-E1162-3

Lot No.
For 200 Assays
2 units (0.005 units / μ L)

Product Description: Human full length C-TAK1, containing a *N*-terminal GST tag, expressed in *E. coli*. Purified by GSH agarose chromatography. The C-TAK1 Positive control is designed to use for Checkpoint Kinase Assay/Inhibitor Screening Kit-1 [Cat# CY-1162]. The C-TAK1 Positive Control should be added to the well at 10 m units/well. For instance, diluted positive control 1:5, use 10 μ L for 1 assay. Unused C-TAK1 Positive control should be stored at -70°C .

Product Size: Full length C-TAK1: 2 units/400 μ L

Formulation: The C-TAK1 Positive Control is supplied frozen in a buffer containing 20mM HEPES-KOH (pH 7.5), 1 % BSA, 1mM EDTA, 2 mM DTT, 50mM NaCl, 0.03 % Brij35 and 50% glycerol.

Source: Human full length C-TAK1, containing *N*-terminal GST tag, expressed in *E. coli*.

Molecular Weight: C-TAK1 demonstrates a double 116 kDa and 72 kDa bands by SDS-PAGE analysis.

Purity: C-TAK1 is greater than 75% pure as determined by SDS-PAGE analysis.

Substrates: C-TAK1 phosphorylates a number of substrates, including PKP2, p53, Cdc25A, Cdc25B and Cdc25C.

Inhibitors: Effective C-TAK1 inhibitor has not been discovered yet.

Unit Definitions: One unit is defined as the amount of kinase required to incorporate 1 nmol of phosphate into the GST-Cdc25C (167-267), per minute at 30°C .

Assay Conditions: Assay activity of C-TAK1 in a 50 μ L reaction containing 20 mM Hepes KOH (pH 7.5), 5 mM MgCl_2 , 1 mM DTT, 100 μ M [γ - ^{32}P] ATP (1 μ Ci), and 4 μ g of GST-Cdc25C fusion protein. Start the reaction by adding 10 μ L of the enzyme, diluted 50-fold in a buffer containing 20 mM Hepes KOH (pH 7.5), 1 mM DTT, 0.03 % Brij35. Incubate for 30 minutes at 30°C . Terminate the reaction by adding 600 μ L of cold 10 % TCA solution containing 0.2 % sodium pyrophosphate and stand on ice for 15 min. Filtrate acid insoluble material through GFC filters (Whatman Inc.), wash 4 times with 1 % TCA and rinse filters with ethanol. Dry filters and count in a liquid scintillation counter.

Storage and Stability: Stable for 12 months at -70°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot enzyme to avoid repeated freezing and thawing.



C-TAK1 Positive Control

Product Data Sheet

For Research Use Only, Not for use in diagnostic procedures

Related Products:

- *Checkpoint Kinase Assay/Inhibitor Screening Kit-1: Cat# CY-1162
- *Chk1 Positive Control: Cat# CY-E1162-1
- *Chk2 Positive Control: Cat# CY-E1162-2

General References:

1. Peng, CY, *et al.*, *Cell Growth Differ.* **9**: 197-208, 1998.
2. Muller J, Ory S, Copeland T, Piwnica-Worms H, Morrison DK. *Mol Cell.* **8**(5):983-93, 2001
3. Kohn EA, Ruth ND, Brown MK, Livingstone M, Eastman A. *J Biol Chem.* **19**;277(29):26553-64, 2002
4. Muller J, Ritt DA, Copeland TD, Morrison DK. *EMBO J.* **22**(17):4431-42, 2003

PRODUCED BY

CycLex Co., Ltd.
1063-103 Terasawaoka
Ina, Nagano 396-0002
Japan
Fax: +81-265-76-7618
e-mail: info@cyclex.co.jp
URL: <http://www.cyclex.co.jp>

CycLex/CircuLex products are supplied for research use only. CycLex/CircuLex products and components thereof may not be resold, modified for resale, or used to manufacture commercial products without prior written approval from CycLex Co., Ltd.. To inquire about licensing for such commercial use, please contact us via email.