



Aurora C Positive Control

Product Data Sheet

For Research Use Only, Not for use in diagnostic procedures

Aurora C Positive Control

(Human, full length, recombinant enzyme expressed in *E. coli*.)

Cat# CY-E1174-2

Lot No.

For 200 Assays

(0.04 units / μL x 200 μL)

Product Description: Human full length Aurora C, containing a *N*-terminal GST-tag, expressed in *E. coli*. Purified by GSH-agarose chromatography. The Aurora C Positive control is designed to use for Aurora Family Assay /Inhibitor Screening Kit [Cat# CY-1174]. The Aurora C Positive Control should be added to the well at 40 m units/well. For instance, diluted positive control 1:10, use 10 μL for 1 assay. Unused Aurora C Positive control should be stored at -70°C .

Product Size: Full length Aurora C: 8 units/200 μL

Formulation: The Aurora C Positive Control is supplied frozen in a buffer containing 20mM HEPES-KOH (pH 7.5), 1 % BSA, 1 mM DTT, 50mM NaCl, 0.03 % Brij35 and 50% glycerol.

Source: Human full length Aurora C, containing *N*-terminal GST-tag, expressed in *E. coli*.

Molecular Weight: Aurora C demonstrates a single 56.4 kDa band by SDS-PAGE analysis.

Purity: Aurora C is greater than 85% pure as determined by SDS-PAGE analysis.

Substrates: Aurora C phosphorylates Histone H3 *in vitro*.

Inhibitors: Specific Aurora C inhibitor has not been discovered yet. However ZM447439 was known as a specific inhibitor for Aurora kinase family ⁽¹⁰⁾.

Unit Definitions: One unit is defined as the amount of kinase required to incorporate 1 nmol of phosphate into the Histone H3, per minute at 30°C .

Assay Conditions: Assay activity of Aurora C in a 50 μL reaction containing 20 mM Hepes KOH (pH 7.5), 5 mM MgCl_2 , 1 mM DTT, 100 μM [γ - ^{32}P] ATP (1 μCi), and 4 μg of Histone H3. Start the reaction by adding 10 μL of the enzyme, diluted 50-fold in a buffer containing 20 mM Hepes KOH (pH 7.5), 1 mM DTT, 0.03 % Brij35. Incubate for 30 minutes at 30°C . Terminate the reaction by adding 600 μL of cold 10 % TCA solution containing 0.2 % sodium pyrophosphate and stand on ice for 15 min. Filtrate acid insoluble material through GFC filters (Whatman Inc.), wash 4 times with 1 % TCA and rinse filters with ethanol. Dry filters and count in a liquid scintillation counter.

Storage and Stability: Stable for 12 months at -70°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot enzyme to avoid repeated freezing and thawing.



Aurora C Positive Control

Product Data Sheet

For Research Use Only, Not for use in diagnostic procedures

Related Products:

- * Aurora A kinase Positive control: Cat# CY-E1165
- * Aurora B kinase Positive control: Cat# CY-E1174-1
- * Aurora-A kinase Assay/Inhibitor Screening Kit: Cat# CY-1165
- * Aurora Family Kinase Assay/Inhibitor Screening Kit: Cat# CY-1174

General References:

1. Francisco, L., Wang, W., and Chan, C.S. (1994). Type 1 protein phosphatase acts in opposition to Ipl1 protein kinase in regulating yeast chromosome segregation. *Mol. Cell Biol.* **14**, 4731–4740.
2. Leveson, J.D., Huang, H.K., Forsburg, S.L., and Hunter, T. (2002). The *Schizosaccharomyces pombe* aurora-related kinase Ark1 interacts with the inner centromere protein Pic1 and mediates chromosome segregation and cytokinesis. *Mol. Biol. Cell* **13**, 1132–1143.
3. Adams, R.R., Carmena, M., and Earnshaw, W.C. (2001a). Chromosomal passengers and the (aurora) ABCs of mitosis. *Trends Cell Biol.* **11**, 49–54.
4. Nigg, E.A. (2001). Mitotic kinases as regulators of cell division and its checkpoints. *Nat. Rev. Mol. Cell Biol.* **2**, 21–32.
5. Kimura, M., Kotani, S., Hattori, T., Sumi, N., Yoshioka, T., Todokoro, K., and Okano, Y. (1997). Cell cycle-dependent expression and spindle pole localization of a novel human protein kinase, Aik, related to Aurora of *Drosophila* and yeast Ipl1. *J. Biol. Chem.* **272**, 13766–13771.
6. Tsai, M.Y., Wiese, C., Cao, K., Martin, O., Donovan, P., Ruderman, J., Prigent, C., and Zheng, Y. (2003). A Ran signalling pathway mediated by the mitotic kinase Aurora A in spindle assembly. *Nat. Cell Biol.* **5**, 242–248.
7. Terada, Y., Tatsuka, M., Suzuki, F., Yasuda, Y., Fujita, S., and Otsu, M. (1998). AIM-1: a mammalian midbody-associated protein required for cytokinesis. *EMBO J.* **17**, 667–676.
8. Kimura, M., Matsuda, Y., Yoshioka, T., and Okano, Y. (1999). Cell cycle-dependent expression and centrosome localization of a third human aurora/Ipl1-related protein kinase, AIK3. *J. Biol. Chem.* **274**, 7334–7340
9. Bischoff, J.R., et al. (1998). A homologue of *Drosophila* aurora kinase is oncogenic and amplified in human colorectal cancers. *EMBO J.* **17**, 3052–3065.
10. Claire Ditchfield, Victoria L. Johnson, Anthony Tighe, Rebecca Ellston, Carolyn Haworth, Trevor Johnson, Andrew Mortlock, Nicholas Keen, and Stephen S. Taylor Aurora B couples chromosome alignment with anaphase by targeting BubR1, Mad2, and Cenp-E to kinetochores. (2003) *J. Cell Biol.* **161**: 267 - 280.

PRODUCED BY

CycLex Co., Ltd.
1063-103 Terasawaoka
Ina, Nagano 396-0002
Japan
Fax: +81-265-76-7618
e-mail: info@cyclex.co.jp
URL: <http://www.cyclex.co.jp>

CycLex/CircuLex products are supplied for research use only. CycLex/CircuLex products and components thereof may not be resold, modified for resale, or used to manufacture commercial products without prior written approval from CycLex Co., Ltd.. To inquire about licensing for such commercial use, please contact us via email.