



Phospho-LSP1 Ser204 Mouse Monoclonal Antibody

Product Data Sheet

For Research Use Only, Not for use in diagnostic procedures

# Phospho-LSP1 Ser204 Mouse Monoclonal Antibody (Clone AT-1E6)

Cat# CY-M1019

100 µg (1 mg/ml x 100 µL)

Clone Name	Applications	Species Cross-reactivity	Molecular Wt.	Source Isotype
AT-1E6	WB, E	H, M, R	52 kDa	Mouse IgG1

**Background:** Signal transduction in B lymphocytes following antigen binding to a membrane-associated immunoglobulin complex initiates a diverse array of intracellular events. To define genes participating in these events, May et al. (1993) and others used various cDNA cloning strategies to identify genes preferentially expressed in B lymphocytes. One gene, variously named LSP1 and pp52, has a number of properties supporting a possible role in B-cell signaling pathways. LSP1 (The leukocyte specific protein 1) is a multi functional protein involved in such divers biological processes as the regulation of neutrophil motility, chemotaxis, adhesion and membrane immunoglobulin M (mIgM) mediated apoptosis of B-lymphocytes. The 330-amino-acid mouse LSP1 protein contains a high-affinity F-actin binding site and in intact cells localizes to the F-actin filament containing cytoskeleton.

**Specificity/Sensitivity:** Phospho-LSP1 Ser204 Antibody detects phosphorylated recombinant LSP1 only when phosphorylated at serine204, by western blotting.

**Source/Purification:** Monoclonal antibody is produced by immunizing mice with a synthetic phosphopeptide corresponding to residues surrounding Ser204 of human LSP1. IgG is purified by protein A-sepharose chromatography.

**Recommended Antibody Dilutions:** Western blotting: 1-2 ug/mL, ELISA for detection of MAPKAP-kinase2 activity: 1 ug/mL

**Storage:** Supplied in 20 mM phosphate buffer (pH 7.5), 300 mM NaCl, 50 % glycerol. Store at -20°C.

**Applications Key:** WB:Western Blotting IP:Immunoprecipitation IHC:Immunohistochemistry IC:Immunocytochemistry F:Flow cytometry E:ELISA FP:Fluorescence Polarization assay

**Species Cross-Reactivity Key:** H:human M:mouse R:rat Hm:hamster Mk:monkey Mi:mink C:chicken X:Xenopus Z:zebra fish All:all species expected Species enclosed in parentheses are predicted to react based on 100% sequence homology.



## Phospho-LSP1 Ser204 Mouse Monoclonal Antibody


### Product Data Sheet

For Research Use Only, Not for use in diagnostic procedures

#### Selected Application References:

1. May, W.; Korenberg, J. R.; Chen, X. N.; Lunsford, L.; Wood, W. J.; Thompson, A.; Wall, R.; Denny, C. T. Human lymphocyte-specific pp52 gene is a member of a highly conserved dispersed family. *Genomics* **15**: 515-520, 1993.
2. Huang CK, Zhan L, Ai Y, Jongstra J. LSP1 is the major substrate for mitogen-activated protein kinase-activated protein kinase 2 in human neutrophils. *J Biol Chem.* **272**(1): 17-9, 1997
3. Jongstra-Bilen J, Janmey PA, Hartwig JH, Galea S, Jongstra J. The lymphocyte-specific protein LSP1 binds to F-actin and to the cytoskeleton through its COOH-terminal basic domain. *J Cell Biol.* **118**(6): 1443-53, 1992
4. Jongstra-Bilen J, Young AJ, Chong R, Jongstra J. Human and mouse LSP1 genes code for highly conserved phosphoproteins. *J Immunol.* **144**(3): 1104-10, 1990

**Fig.1 ELISA for measurement of recombinant MAPKAP-kinase2 activity using Phospho-LSP1 Ser204 Monoclonal antibody (AT-1E6) in CycLex® MAPKAP-kinase 2 Assay/Inhibitor Screening Kit (Cat# CY-1166)**



#### Western Immunoblotting Protocol

##### Solutions and Reagents

Note: Prepare solutions with Milli-Q or equivalently purified water.

**Transfer Buffer:** 25 mM Tris base, 0.2 M glycine, 20% methanol (pH 8.5)

**SDS Sample Buffer (1X):** 62.5 mM Tris-HCl (pH 6.8 at 25°C), 2% w/v SDS, 10% glycerol, 50 mM DTT, 0.01% w/v bromophenol blue or phenol red

**Blocking Buffer:** 1X TBS, 0.1% Tween-20 with 5% w/v nonfat dry milk; for 150 ml, add 15 ml 10X TBS to 135 ml water, mix. Add 7.5 g nonfat dry milk and mix well. While stirring, add 0.15 ml Tween-20 (100%).

**10X TBS (Tris-buffered saline):** To prepare 1 liter of 10X TBS: 24.2 g Tris base, 80 g NaCl; adjust pH to 7.6 with HCl (use at 1X).

**Primary Antibody Dilution Buffer:** 1X TBS, 0.1% Tween-20 with 5% blocking agent; for 20 ml, add 2 ml 10X TBS to 18 ml water, mix. Add 1.0 g BSA and mix well. While stirring, add 20 µl Tween-20



## Phospho-LSP1 Ser204 Mouse Monoclonal Antibody

### Product Data Sheet

**For Research Use Only, Not for use in diagnostic procedures**

(100%).

**Chemiluminescent HRP Detection:** secondary anti-rabbit antibody conjugated to horseradish peroxidase (HRP), ECL™ chemiluminescent reagent (Amersham Pharmacia)

**Wash Buffer TBS/T:** 1X TBS, 0.1% Tween-20

**Blotting Membrane:** This protocol has been optimized for nitrocellulose membranes, which we recommend. PVDF membranes may also be used.

#### Protein Blotting

A general protocol for sample preparation is described below.

1. Treat cells by adding fresh media containing regulator for desired time.
2. Aspirate media from cultures; wash cells with 1X PBS; aspirate.
3. Lyse cells by adding 1X SDS Sample Buffer (100  $\mu$ l per well of 6-well plate or 500  $\mu$ l per plate of 10 cm<sup>2</sup> plate). Immediately scrape the cells off the plate and transfer the extract to a microcentrifuge tube. Keep on ice.
4. Sonicate for 10–15 seconds to shear DNA and reduce sample viscosity.
5. Heat a 20  $\mu$ l sample to 95–100°C for 5 minutes; cool on ice.
6. Microcentrifuge for 5 minutes.
7. Load 20  $\mu$ l onto SDS-PAGE gel (10 cm x 10 cm).
8. Electrotransfer to nitrocellulose membrane.

#### Membrane Blocking and Antibody Incubations

*Note: Volumes are for 10 cm x 10 cm (100 cm<sup>2</sup>) of membrane; for different sized membranes, adjust volumes accordingly.*

1. (Optional) After transfer, wash nitrocellulose membrane with 25 ml TBS for 5 minutes at room temperature.
2. Incubate membrane in 25 ml of Blocking Buffer for 1 hour at room temperature.
3. Wash 3 times for 5 minutes each with 15 ml of TBS/T.
4. Incubate membrane and primary antibody (at the appropriate dilution) in 10 ml Primary Antibody Dilution Buffer with gentle agitation overnight at 4°C.
5. Wash 3 times for 5 minutes each with 15 ml of TBS/T.
6. Incubate membrane with HRP-conjugated secondary antibody (1:3000 in 10 ml of Blocking Buffer with gentle agitation for 1 hour at room temperature.
7. Wash 3 times for 5 minutes each with 15 ml of TBS/T.

#### Detection of Proteins

1. Incubate membrane with 4 ml ECL™ with gentle agitation for 1 minute at room temperature.
2. Drain membrane of excess developing solution, do not let dry, wrap in plastic wrap and expose to x-ray film. An initial ten seconds exposure should indicate the proper exposure time.

#### Immunoprecipitation Followed by Western Immunoblotting Protocol

##### Solutions and Reagents

*Note: Prepare solutions with Milli-Q or equivalently purified water.*

**Cell Lysis Buffer (1X):** 20 mM Tris (pH 7.5), 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% Triton X-100, 2.5 mM sodium pyrophosphate, 1 mM Glycerolphosphate, 1 mM Na<sub>3</sub>VO<sub>4</sub>, 1  $\mu$ g/ml Leupeptin

*Note: We recommend adding 1 mM PMSF before use.*

**Protein A Agarose Beads:** Add 5 ml of 1X PBS to 1.5 g of Protein A Agarose Beads. Shake 2 hours at 4°C; spin down. Wash pellet twice with PBS. Resuspend beads in 1 volume of PBS. (Can be stored for 2 weeks at 4°C)

**3X SDS Sample Buffer:** 187.5 mM Tris-HCl (pH 6.8 at 25°C), 6% w/v SDS, 30% glycerol, 150 mM



## Phospho-LSP1 Ser204 Mouse Monoclonal Antibody

### Product Data Sheet

**For Research Use Only, Not for use in diagnostic procedures**

DTT, 0.03% w/v bromophenol blue,

**Transfer Buffer:** 25 mM Tris base, 0.2 M glycine, 20% methanol (pH 8.5)

**Blocking Buffer:** 1X TBS, 0.1% Tween-20 with 5% w/v nonfat dry milk. For 150 ml, add 15 ml 10X TBS to 135 ml water, mix. Add 7.5 g nonfat dry milk and mix well. While stirring, add 0.15 ml Tween-20 (100%).

**10X TBS (Tris-buffered saline):** For 1 liter of 10X TBS, use 24.2 g Tris base and 80 g NaCl. Adjust pH to 7.6 with HCl (use at 1X).

**Primary Antibody Dilution Buffer:** 1X TBS, 0.05% Tween-20 with 5% nonfat dry milk. For 20 ml, add 2 ml 10X TBS to 18 ml water, mix. Add 1.0 g nonfat dry milk and mix well. While stirring, add 10  $\mu$ l Tween-20 (100%).

**Wash Buffer TBS/T:** 1X TBS, 0.1% Tween-20

**Chemiluminescent HRP Detection:** secondary anti-rabbit antibody conjugated to horseradish peroxidase (HRP), ECL<sup>TM</sup> chemiluminescent reagent (Amersham Pharmacia)

**Wash Buffer TBS/T:** 1X TBS, 0.1% Tween-20

**Blotting Membrane:** This protocol has been optimized for nitrocellulose membranes, which we recommend. PVDF membranes may also be used.

#### Preparing Cell Lysates

1. Aspirate media. Treat cells by adding fresh media containing regulator for desired time.
2. To harvest cells under non-denaturing conditions, remove media and rinse cells once with ice-cold PBS.
3. Remove PBS and add 0.5 ml 1X ice-cold Cell Lysis Buffer plus 1 mM PMSF to each plate (10 cm<sup>2</sup>) and incubate the plate on ice for 5 minutes.
4. Scrape cells off the plate and transfer to microcentrifuge tubes. Keep on ice.
5. Sonicate 4 times for 5 seconds each on ice.
6. Microcentrifuge for 10 minutes at 4°C, and transfer the supernatant to a new tube. The supernatant is the cell lysate. If necessary, lysate can be stored at -80°C.

#### Immunoprecipitation

1. Take 200  $\mu$ L cell lysate and add primary antibody; incubate with gentle rocking overnight at 4°C.
2. Add Protein A Agarose Beads (20  $\mu$ L of 50% bead slurry). Incubate with gentle rocking for 1-3 hours at 4°C.
3. Microcentrifuge for 30 seconds at 4°C. Wash pellet 2 times with 500  $\mu$ L of 1X Cell Lysis Buffer. Keep on ice during washes.
4. Resuspend the pellet with 20  $\mu$ L 3X SDS Sample Buffer. Vortex, then, microcentrifuge for 30 seconds.
5. Heat the sample to 95-100°C for 2-5 minutes.
6. Load the sample (15-30  $\mu$ L) on SDS-PAGE gel (12-15%).
7. Analyze sample by Western blotting (see Western Immunoblotting Protocol).

#### PRODUCED BY

CycLex Co., Ltd.

1063-103 Terasawaoka

Ina, Nagano 396-0002

Japan

Fax: +81-265-76-7618

e-mail: [info@cyclex.co.jp](mailto:info@cyclex.co.jp)

URL: <http://www.cyclex.co.jp>

CycLex/CircuLex products are supplied for research use only. CycLex/CircuLex products and components thereof may not be resold, modified for resale, or used to manufacture commercial products without prior written approval from CycLex Co., Ltd.. To inquire about licensing for such commercial use, please contact us via email.