

## p38 delta (Human), Active

### Full-length recombinant protein expressed in Sf9 cells

Cat# CY-SPM37

Lot No. K145-2  
5 µg 0.1 µg/µl

#### Background:

p38  $\delta$  (p38-delta, SAPK4) is a member of the p38 MAPK family and is activated by chemical and environmental stresses as well as by proinflammatory cytokines. p38  $\delta$  has a TGY dual phosphorylation motif and is activated in response to cellular stresses and proinflammatory cytokines (1). MAP kinase kinases 3, and 6 can phosphorylate and activate this kinase. Transcription factor ATF2, and microtubule dynamics regulator stathmin have been shown to be the substrates of this kinase (2).

#### Product Description:

Recombinant full-length human p38  $\delta$  was expressed by baculovirus in Sf9 insect cells using a N-terminal GST tag. The gene accession number is NM\_002754.

#### Gene Aliases:


SAPK4; PRKM13; MAPK13

#### Formulation:

Recombinant protein stored in 50mM Tris-HCl, pH 7.5, 150mM NaCl, 0.25mM DTT, 0.1mM EGTA, 0.1mM EDTA, 0.1mM PMSF, 25% glycerol.

#### Purity & Molecular Weight:

The purity was determined to be >80% by densitometry. Approx. MW 71kDa.



#### Storage:


Store product at  $-70^{\circ}\text{C}$ . For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

#### Stability:

Unopened vial at  $-70^{\circ}\text{C}$ , 1 year from date of shipment.

**Specific Activity:**

The specific activity was determined to be 259 nmol/min/mg as per Activity Assay Protocol.

**Activity Assay Protocol:**

Assay activity of the kinase in a 25  $\mu$ L reaction consisting of 5  $\mu$ L of 5 X Kinase Assay Buffer, 10  $\mu$ L of 1 mg/ml the Substrate Solution, 5  $\mu$ L of diluted kinase and 5  $\mu$ L of 250  $\mu$ M ATP solution containing [ $\gamma$ - $^{32}$ P] ATP (0.167  $\mu$ Ci/ $\mu$ L). Start the reaction by adding the ATP solution. Incubate for 15 minutes at 30°C. Terminate the reaction by spotting 20  $\mu$ L of the reaction mixture onto phosphocellulose P81 paper. Air-dry the P81 paper and sequentially wash 4 times for approximately 10 minutes each in 1% phosphoric acid with constant gentle stirring. Count the P81 paper in a liquid scintillation counter.

**Substrate Solution:**

Myelin basic protein (MBP) diluted in distilled H<sub>2</sub>O to a final concentration of 1 mg/ml.

**5 X Kinase Assay Buffer:**

25mM MOPS, 12.5mM  $\beta$ -glycerol-phosphate, 25mM MgCl<sub>2</sub>, 5mM EGTA, 2mM EDTA. Add 0.25mM DTT to Kinase Assay Buffer prior to use.

**References:**

1. Goedert, M. et al: Activation of the novel stress-activated protein kinase SAPK4 by cytokines and cellular stresses is mediated by SKK3 (MKK6); comparison of its substrate specificity with that of other SAP kinases. EMBO J, 16: 3563-3571, 1997.
2. Kumar, S. et al: Novel homologues of CSBP/p38 MAP kinase: activation, substrate specificity and sensitivity to inhibition by pyridinyl imidazoles. Biochem Biophys Res Commun. 1997 Jun 27;235(3):533-8.

**CycLex Co., Ltd**

**1063-103 Tera-Sawaoka, Ina, Nagano, Japan 396-0002**

**Fax: 81-265-76-7618**

**e-mail: [info@cyclex.co.jp](mailto:info@cyclex.co.jp)**

**URL: <http://www.cyclex.co.jp>**