

## SRC (Rous sarcoma virus), Active

### Full-length recombinant protein expressed in Sf9 cells

Cat# CY-SPS19-V

Lot No. B176-1  
5 µg 0.1 µg/µl

#### Background:

SRC family belongs to non-receptor tyrosine kinases. SRC was originally identified as a transforming protein of the Rous sarcoma virus (RSV) that had enzymatic ability to phosphorylate tyrosine in protein substrates (1). SRC is overexpressed and activated in a large number of human malignancies and has been linked to the development of cancer and progression to distant metastases (2). In addition to increasing cell proliferation, a key role of SRC in cancer seems to be the ability to promote invasion and motility, functions that might contribute to tumour progression.

#### Product Description:

Recombinant full-length Rous sarcoma virus SRC was expressed by baculovirus in Sf9 insect cells using a N-terminal GST tag. The gene accession number is M11753.

#### Gene Aliases:


ASV; SRC1; p60-Src

#### Formulation:

Recombinant protein stored in 50mM Tris-HCl, pH 7.5, 150mM NaCl, 0.25mM DTT, 0.1mM EGTA, 0.1mM EDTA, 0.1mM PMSF, 25% glycerol.

#### Purity & Molecular Weight:

The purity was determined to be >90% by densitometry. Approx. MW 85kDa.



#### Storage:


Store product at  $-70^{\circ}\text{C}$ . For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

#### Stability:

Unopened vial at  $-70^{\circ}\text{C}$ , 1 year from date of shipment.

**Specific Activity:**

The specific activity was determined to be 116 nmol/min/mg as per Activity Assay Protocol.



**Activity Assay Protocol:**

Assay activity of the kinase in a 25  $\mu$ L reaction consisting of 5  $\mu$ L of 5 X Kinase Assay Buffer, 10  $\mu$ L of 1 mg/ml the Substrate Solution, 5  $\mu$ L of diluted kinase and 5  $\mu$ L of 250  $\mu$ M ATP solution containing [ $\gamma$   $^{32}$ P] ATP (0.167  $\mu$ Ci/ $\mu$ L). Start the reaction by adding the ATP solution. Incubate for 15 minutes at 30°C. Terminate the reaction by spotting 20  $\mu$ L of the reaction mixture onto phosphocellulose P81 paper. Air-dry the P81 paper and sequentially wash 4 times for approximately 10 minutes each in 1% phosphoric acid with constant gentle stirring. Count the P81 paper in a liquid scintillation counter.

**Substrate Solution:**

SRC synthetic peptide substrate (KVEKIGEGTYGVVYK) diluted in distilled H<sub>2</sub>O to a final concentration of 1 mg/ml.

**5 X Kinase Assay Buffer:**

25mM MOPS, 12.5mM  $\beta$ -glycerol-phosphate, 20mM MgCl<sub>2</sub>, 25mM MnCl<sub>2</sub>, 5mM EGTA, 2mM EDTA. Add 0.25mM DTT to Kinase Assay Buffer prior to use.

**References:**

1. Collett, M S. et al: Protein kinase activity associated with the avian sarcoma virus src gene product. Proc Natl Acad Sci U S A. 1978 Apr;75(4):2021-4.
2. Jacobs, C. et al: Expression of pp60c-src protein kinase in adult and fetal human tissue: high activities in some sarcomas and mammary carcinomas. Cancer Res. 1983 Apr;43(4):1696-702.

**CycLex Co., Ltd**

**1063-103 Tera-Sawaoka, Ina, Nagano, Japan 396-0002**

**Fax: 81-265-76-7618**

**e-mail: [info@cyclex.co.jp](mailto:info@cyclex.co.jp)**

**URL: <http://www.cyclex.co.jp>**